

Reliable on-hand treatment for episodes of frequent seizure activity

For pharmacists looking to place orders for VALTOCO, here's what you need to know.



Please contact your wholesaler to place orders for VALTOCO

Dose	NDC Number	What's Included	Pkg. Size/ Pkg. Quantity	Box Dimensions/ Weight
5 mg	72252-505-05	This box contains five (5) individual blister packs. Each blister pack contains 1 VALTOCO nasal spray device. Each VALTOCO nasal spray device contains 5 mg diazepam in 0.1 mL solution. The entire carton is to be dispensed as a unit.	1/5	3.473" x 2.661" x 5.903" / 0.189 lb
10 mg	72252-510-05	This box contains five (5) individual blister packs. Each blister pack contains 1 VALTOCO nasal spray device. Each VALTOCO nasal spray device contains 10 mg diazepam in 0.1 mL solution. The entire carton is to be dispensed as a unit.	1/5	3.473" x 2.661" x 5.903" / 0.189 lb
15 mg	72252-515-10	This box contains five (5) individual blister packs. Each blister pack contains 2 VALTOCO nasal spray devices. Each VALTOCO nasal spray device contains 75 mg diazepam in 0.1 mL solution. Use both nasal spray devices for the full 15 mg dose. The entire carton is to be dispensed as a unit.	1/5	4.223" x 3.473" x 5.903" / 0.29 lb
20 mg	72252-520-10	This box contains five (5) individual blister packs. Each blister pack contains 2 VALTOCO nasal spray devices. Each VALTOCO nasal spray device contains 10 mg diazepam in 0.1 mL solution. Use both nasal spray devices for the full 20 mg dose. The entire carton is to be dispensed as a unit.	1/5	4.223" x 3.473" x 5.903" / 0.29 lb

Store at room temperature between $68^{\circ}F$ and $77^{\circ}F$ ($20^{\circ}C$ to $25^{\circ}C$). Do not freeze. Protect from light.¹

VALTOCO is indicated for the acute treatment of intermittent, stereotypic episodes of frequent seizure activity (ie, seizure clusters, acute repetitive seizures) that are distinct from a patient's usual seizure pattern in patients with epilepsy 6 years of age and older.

Please see complete Instructions for Use on **VALTOCOHCP.com**.

How to administer VALTOCO²

Remove VALTOCO from the blister pack.



HOLD VALTOCO with your thumb on the bottom of the plunger and your first and middle fingers on either side of the nozzle.

DO NOT test or prime; each device sprays only one time.



INSERT the tip of the nozzle into 1 nostril until your fingers, on either side of the nozzle, are against the bottom of the person's nose.

For nasal use only.



PRESS the bottom of the plunger firmly with your thumb to give VALTOCO.

Throw away nasal spray device(s) after use.

If giving the **15 mg or 20 mg dose**, repeat the steps using the second device in the other nostril to give the full dose of VALTOCO.

Remember: If needed, a second dose may be given at least 4 hours after initial dose.

Patients should not use more than 2 doses of VALTOCO to treat a single episode.

These are not the full Instructions for Use. Please see the complete

Instructions for Use at VALTOCOHCP.com.



Frequently asked questions about VALTOCO



1. How frequently can VALTOCO be dosed?

It's recommended that VALTOCO be used to treat no more than 1 episode every 5 days and no more than 5 episodes per month. If needed, a second dose may be given at least 4 hours after initial dose. Patients should not use more than 2 doses of VALTOCO to treat a single episode.¹

2. What is the most important information I should know about VALTOCO?

VALTOCO is a benzodiazepine medicine. Benzodiazepines have warnings for risks from concomitant use with opioids; abuse, misuse, and addiction; and dependence and withdrawal reactions.¹

See below for additional important safety information.

3. Does VALTOCO need to be primed?

No. Patients or their care partners SHOULD NOT test or prime VALTOCO. Each device sprays only one time.²

4. Can patients carry VALTOCO with them?

Yes. With the small, portable, and discreet packaging of VALTOCO, patients can carry it with them in their backpack or purse—whenever, wherever. It does not need to be refrigerated and is designed for prompt administration by anyone.¹

5. How can I get more information about VALTOCO?

To learn more about VALTOCO, or to request a visit from your local Neurelis representative, go to **VALTOCOHCP.com**.

Indication

VALTOCO® (diazepam nasal spray) is indicated for the acute treatment of intermittent, stereotypic episodes of frequent seizure activity (ie, seizure clusters, acute repetitive seizures) that are distinct from a patient's usual seizure pattern in patients with epilepsy 6 years of age and older.

IMPORTANT SAFETY INFORMATION

WARNING: RISKS FROM CONCOMITANT USE WITH OPIOIDS; ABUSE, MISUSE, AND ADDICTION; and DEPENDENCE AND WITHDRAWAL REACTIONS

- Concomitant use of benzodiazepines and opioids may result in profound sedation, respiratory depression, coma, and death.
- The use of benzodiazepines, including VALTOCO, exposes users to risks of abuse, misuse, and addiction, which can lead to overdose or death.
- The continued use of benzodiazepines may lead to clinically significant physical dependence. Although VALTOCO is indicated only for intermittent use, if used more frequently than recommended, abrupt discontinuation or rapid dosage reduction of VALTOCO may precipitate acute withdrawal reactions, which can be life-threatening.

Adverse Reactions

The most common adverse reactions (at least 4%) were somnolence, headache, and nasal discomfort.

Diazepam, the active ingredient in VALTOCO, is a Schedule IV controlled substance.

To report SUSPECTED ADVERSE REACTIONS, contact Neurelis, Inc. at 1-866-696-3873 or FDA at 1-800-FDA-1088 (www.fda.gov/medwatch).

Please see full Prescribing Information, including Boxed Warning.

References: 1. Valtoco. Prescribing Information. Neurelis Inc. https://www.valtoco.com/PI; **2.** Valtoco. Instructions for Use. Neurelis Inc. https://www.valtoco.com/sites/default/files/pdf/Instructions_For_Use.pdf

